

*PERFORMANCE OF THE AIRBORNE ULTRA HIGH SENSITIVITY
AEROSOL SPECTROMETER DURING THE CARBONACEOUS
AEROSOL AND RADIATIVE EFFECTS STUDY AND CALWATER
FIELD CAMPAIGNS*

Jason Tomlinson, John Shilling, Mikhail Pekour, Rahul Zaveri, Jennifer
Comstock, John Hubbe, Duli Chand, Celine Kluzek, and Beat Schmid,
Pacific Northwest National Laboratory, WA 99352

Haflidi Jonsson and Roy Woods
Center for Interdisciplinary Remotely Piloted Aircraft Studies, CA, 93933

Jian Wang, Stephen Springston, Chongai Kuang, and Lawrence Kleinman
Brookhaven National Laboratory, NY, 11973

Kimberly Prather, Kaitlyn Suski, and John Cahill
University of San Diego, CA, 92093

For presentation at the
30th Annual Conference of the American Association for Aerosol Research,
Orlando, FL
Oct. 3-7, 2011

**Environmental Sciences Department/Atmospheric Sciences Division
Brookhaven National Laboratory**
P.O. Box, Upton, NY
www.bnl.gov

ABSTRACT

Anthropogenic aerosols effect on the formation of clouds and precipitation amounts is one of the largest uncertainties in the prediction of global climate change. To determine the magnitude of this effect, the Airborne Ultra High Sensitivity Aerosol Spectrometer (UHSAS) is being used to measure aerosol size distributions from 0.060 to 1.0 mm at a frequency of 1HZ. This high temporal resolution enables detailed analysis of urban pollution plumes and the aerosol in the proximity of clouds. The instrument was recently operated aboard the Department of Energy (DOE) Atmospheric Radiation Measurement (ARM) Aerial Facility Gulfstream-1 aircraft during the Carbonaceous Aerosol and Radiative Effects Study in June 2010 and the CalWater field campaign in February 2011. The measured size distributions will be compared with the measured size distributions from other instruments operated aboard the aircraft to determine the overall performance of the instrument within different environmental conditions and when sampling different mixtures of aerosol composition. In addition, a preliminary analysis of the aerosol and the aerosol effect on clouds over the California Central Valley and the Sierra Nevada mountain range during the summer season, winter season, atmospheric river events, and long-range transport events from Asia will be presented.

NOTICE: This manuscript has been authored by employees of Brookhaven Science Associates, LLC under Contract No. DE-AC02-98CH10886 with the U.S. Department of Energy. The publisher by accepting the manuscript for publication acknowledges that the United States Government retains a non-exclusive, paid-up, irrevocable, world-wide license to publish or reproduce the published form of this manuscript, or allow others to do so, for United States Government purposes.